

Project Title: *Robotic Therapy for Force Training of the Upper Extremity in Chronic Hemiparetic Stroke*

Investigators: Leonard Kahn (co-PI), David Reinkensmeyer (co-PI), W. Zev Rymer (co-investigator), Heidi Waldinger, Heidi Roth, Colby Smith

Project description

The goal of this project is to use a novel robotic therapy device, the ARM Guide, to improve recovery of upper extremity function after stroke. We have already demonstrated in an initial pilot study that the device can improve movement recovery following chronic stroke. However, the results from our study, combined with evidence from other recent studies of robotic therapy, suggest that robotic therapy might be substantially enhanced by specifically targeting abnormal force generation after stroke, a therapeutic paradigm in which subjects will use biofeedback of arm forces to consciously direct their movements towards a desired target. The current project implements this novel form of robotic therapy, called “guided force training”, and tests whether this therapy can improve recovery beyond previous forms of robotic and conventional therapy.

We are examining 72 stroke survivors who are at least six months post-injury with unilateral lesion. Subjects participate in one of three therapy programs across 24 one-hour sessions in eight weeks. The first program, free reaching therapy, involves subjects practicing unsupported reaching movements with the affected arm to an array of targets throughout the workspace. A second group participates in conventional therapy during one-on-one sessions with an occupational therapist. The strategies employed by the therapist concentrate on scapular and shoulder alignment, passive and active assisted range of motion, and reeducation using a task-oriented approach. Finally, the robotic therapy group engages in reaching therapy with the device. As the subject attempts to make reaching movements while holding onto a handpiece along a linear track (see figure at right), a force sensor in the handpiece records the direction of forces generated by the user. If impaired coordination causes the user to push the handpiece in a direction other than towards the intended target, the motor will halt movement and the user will be given graphical feedback showing them how to correct their reach. This punishment-reward system (proper endpoint force rewarded by allowing smooth, unimpeded movement) trains users to consciously aim hand movements toward the desired target.



Progress to date

We have enrolled fifteen subjects and ten have completed the training program – four in free reaching, three in conventional therapy, and three in guided force training. Current results show positive changes in upper extremity function during simple reaching movements and more complex activities of daily living. However, our participant numbers are still small and no differences have yet been found between groups to draw any conclusions about the relative efficacies of these therapies.